

Storage of pollen and properties of olive stigma for breeding purposes

Armazenamento de pólen e propriedades do estigma de oliveira para fins de melhoramento genético

ABSTRACT - In order to ensure success in controlled hybridizations in olive tree cultivation, the information on pollen viability and stigma receptivity is essential. The aim was to establish methodologies that increase the preservation of pollen viability and to establish the time to perform crossbreeds in hybridization studies with olive trees. Three experiments were performed with plants from the cultivar Arbequina, in Maria da Fé, MG, Brazil. In the first experiment, the description of the flower events was performed. In the second, anthers were desiccated in eppendorfs, being stored at three different conditions for pollen viability test: room temperature (27 °C), refrigerator (8 °C) and freezer (-10 °C). In order to evaluate the *in vitro* germination, culture medium for olive pollen grains was used. In this respect, pollen grains were transferred in Petri dish containing culture medium and placed in a BOD (biochemical oxygen demand) chamber at 28 °C for 60 h, being counted. The first evaluation was performed prior to the assembly of the experiment, testing the initial viability, whereas the second occurred 24 h after storage. Subsequently, seven evaluations were performed fortnightly. In the third experiment, the stigma receptivity was verified by the 3% hydrogen peroxide method, with flowers in pre-anthesis, anthesis and post-anthesis, evaluated hourly in the period from 7 a.m. to 6 p.m. for three days. In the description of the flower events, it was verified that the olive tree shows diurnal

22 anthesis, with flower opening between 10 a.m. and 11 a.m. The anthers stored in a freezer
23 preserved the viability for 60 days and the stigmas were receptive since the pre-anthesis.

24 **Key words:** *Olea europaea* L.. Pollen viability. Pollen grains. Hybridization.

25 **RESUMO** - Para assegurar o sucesso em hibridizações controladas na cultura da oliveira, é
26 fundamental informações sobre a viabilidade polínica, assim como a receptividade do estigma.
27 Objetivou-se estabelecer metodologias que aumentem a preservação da viabilidade polínica e
28 estabelecer o momento para se realizar cruzamentos em trabalhos de hibridização com oliveiras.
29 Foram realizados três experimentos com plantas da cultivar Arbequina, em Maria da Fé, MG. No
30 primeiro experimento realizou-se a descrição dos eventos florais. No segundo, anteras foram
31 dessecadas em eppendorfs, sendo armazenados em três condições diferentes para teste de
32 viabilidade polínica : temperatura ambiente (27 °C), geladeira (8 °C) e freezer (-10 °C). Para
33 avaliação da germinação *in vitro*, utilizou-se meio de cultura para grãos de pólen de oliveira. Para
34 isso, grãos de pólen foram transferidos em placa de Petri, contendo meio de cultura e colocadas
35 em estufa do tipo BOD (Demanda Biológica de Oxigenio) a 28 °C por 60 h, sendo
36 contabilizados. A primeira avaliação foi realizada antes da montagem do experimento, testando a
37 viabilidade inicial: a segunda 24 horas após armazenamento. Posteriormente, foram realizadas
38 sete avaliações quinzenais. No terceiro experimento verificou-se a receptividade estigmática pelo
39 método de peróxido de hidrogênio 3%, com flores em pré-antese, antese e pós-antese, avaliados a
40 cada hora, no período de 7 h as 18 h por três dias. Na descrição dos eventos florais, verificou-se
41 que a oliveira apresenta antese diurna, com abertura floral entre 10 e 11 horas. As anteras
42 armazenadas em freezer preservam a viabilidade por 60 dias e os estigmas apresentaram-se
43 receptivos desde a pré-antese.

44 **Palavras-chave:** *Olea europaea* L.. Viabilidade polínica. Grãos de pólen. Hibridação.

45

46

INTRODUCTION

47 The olive tree (*Olea europaea* L.) is characterized as an allogamous plant with self-
48 incompatibility of pollen-pistil or gamete, being observed an abundant flowering and a low
49 fertilization (lower than 20% of the flowers are differentiated into fruits) (KOUBOURIS;
50 METZIDAKIS; VASILAKAKIS, 2009). Its inflorescences are constituted by actinomorphic
51 flowers with regular symmetry, composed of four sepals, four petals, two stamens and one
52 central pistil (OLIVEIRA, M., *et al.*, 2012).

53 Brazil is the eighth largest importer of olive oil and table olive in the world (COI, 2017).
54 Research on olive groves has been performed in some regions of the states of Rio Grande do Sul
55 and Minas Gerais (SILVA *et al.*, 2012a).

56 The carpometric trait of fruits and olive pits is decisive to define the exploitation potential
57 of the cultivar, either highlighting its purpose of usage, for olive oil extraction or canning. In this
58 respect; some cultivars are highlighted, such as Negroa, MGS ASC322, MSG MISS293, MGS
59 JB1, MGS GRAP556, MGS GRAP541 and Arbequina (SILVA *et al.*, 2012b).

60 Arbequina cultivar is the most commonly used by the producers and hence is the most
61 representative cultivar in the Brazilian olive groves among the cultivars with productive potential
62 evaluated in the South and Southern regions (SILVA *et al.*, 2016). It is a cultivar originating from
63 Spain used to produce olive oil, which was very well adapted to the environmental conditions of
64 the South and Southeast of Brazil (SILVA *et al.*, 2012a).

65 Currently, the productive performance of different cultivars introduced in the recent years
66 in the country is already known (SILVA *et al.*, 2012b). Brazil also has varieties developed from
67 Brazilian breeding programs, whose study was performed based on natural hybridizations, with
68 subsequent evaluation of the performance of genotypes from these crossbreeds (OLIVEIRA, A.,
69 *et al.*, 2012). However, the new premise for breeding studies would be the use of controlled
70 hybridizations, in which information would be obtained from both parents, increasing the
71 possibility of finding a cultivar with the desired characteristics within a selection study.

72 To ensure success in hybridizations, the pollen should essentially show high rate of
73 viability and germination. Theoretically, the pollen collection in the proper and correctly
74 prepared flower stage does not require a viability test (BRITO *et al.*, 2010).

75 However, there is often no possibility of immediate use, such as in cases where there is a
76 lack of flower synchronicity among the cultivars, or when the pollen grains are collected in a
77 region far from that of hybridization. In these situations, it is necessary to evaluate the viability
78 before using the material, as well as to test and develop techniques that extend its viability
79 (MOURA; MACHADO; LÈDO, 2015).

80 The reproductive structure denominated stigma is another fundamental factor, since the
81 aptitude of flowers in the fertilization process is directly related to its receptivity to the pollen
82 grain. Such receptivity can last only a few hours, such as in some species of the family
83 Anacardiaceae, or several days as in several species of the family Solanaceae (SCHIFINO-
84 WITTMANN; DALL'AGNOL, 2002). In some plants, the ovary receptivity is indicated by the
85 moisture of the estimate, which allows adhering the pollen grain. (SCHIFINO-WITTMANN;
86 DALL'AGNOL, 2002).

87 Therefore, the aim was to establish methodologies that preserve pollen viability over time
88 and to determine the moment of stigmatic receptivity of olive flowers to perform crossbreeds in
89 hybridization studies.

90 MATERIAL AND METHODS

91 The present study was performed with the cultivar Arbequina, aged eight years, in Maria da
92 Fé, MG, Brazil (22°18'51" S, 45°23'24" W and 1,276 m altitude) between August and September
93 2015 (SILVA *et al.*, 2016).

94 **1st experiment: description and flower events**

95 For the determination of anthesis, daily quantitative observations of 50 flower buds from 10
96 randomly selected individuals were performed and noted from 7 a.m. to 6 p.m. for 20 days
97 (BRITO *et al.*, 2010). In this phase, the main flower opening time, flower synchronicity (within
98 an inflorescence and among different inflorescences), onset of anthesis and flower longevity were
99 observed. Results were expressed as a percentage.

100 **2nd experiment: conservation of pollen grains**

101 In this second phase, 15 plants were selected, being collected randomly 560 flower buds at
102 "balloon" stage, in pre-anthesis in the morning.

103 In the laboratory, the anthers were separated with the aid of a forceps, being placed four
104 anthers by eppendorfs, totaling 280 eppendorfs. Afterwards, the tubes were desiccated with
105 silica gel and maintained at 28 °C for 24 h, so that the anthesis occurred and the pollen grains had
106 their moisture content reduced (CUCHIARA; SILVA; BOBROWSHI, 2012). Then, the test was
107 performed to determine the initial viability of pollen grains, using 10 eppendorfs, being one per
108 Petri dish, using culture medium specific for olive pollen germination (SILVA *et al.*, 2016).

109 Subsequently, the remainder of eppendorfs were distributed among the treatments (90
110 epperddorfs per treatment).

111 The used treatments consisted of keeping the pollen grains stored under three different
112 conditions for later viability test: room temperature (27 °C), refrigerator (8 °C) and freezer (-10
113 °C). After 24 h, a new evaluation was performed, followed by seven evaluations fortnightly. A
114 total of 10 eppendorfs per treatment (replicates) were tested in each evaluation period.

115 The experimental design was completely randomized in a 9 x 3 factorial design (storage
116 time x environments) and 10 replicates, each replication consisting of one Petri dish, with the
117 content of 1 eppendorf, where germinated and non-germinated pollen were counted in five
118 randomly selected fields of view

119 In order to determine the viability of pollen grains, the *in vitro* germination method was
120 used through culture medium specific for germination of olive pollen grains. The culture medium
121 consisted of 4 g L⁻¹ agar plus 90 g L⁻¹ sucrose, 400 mg L⁻¹ of boric acid and pH adjusted to 5.79,
122 in the absence of calcium nitrate and magnesium sulfate, kept for 60 h at 28 °C (SILVA *et al.*,
123 2016).

124 For each stage, the pollen grains were transferred with the aid of a brush to a Petri dish
125 surface containing 20 mL of culture medium. Subsequently, the Petri dishes were kept in
126 germination chamber (BOD) in the absence of light at 28°C for 60 h, being counted germinated
127 and ungerminated pollen grains using a microscope with 10x objective lens (SILVA *et al.*, 2016).

128 The pollen grain, whose length of the pollen tube exceeded twice the diameter was
129 considered as germinated (FIGUEIREDO *et al.*, 2013). This experiment was performed in a

130 completely randomized design with three replicates, being each replicate consisting of one
131 eppendorfs and five fields of view per Petri dish.

132 **3rd experiment: receptivity of stigma**

133 Stigma receptivity was tested using the 3% hydrogen peroxide (H₂O₂) method. This method
134 consists of the deposition of 3% hydrogen peroxide on the stigma of flowers, in which the
135 stigmas that showed bubble formation were considered as receptive (KEARNS; INOUYE, 1993).

136 For this experiment, a completely randomized design was used, and the evaluations were
137 performed from 7 a.m. to 6 p.m. for three days, with 10 replicates. Stigma receptivity was
138 evaluated in three flower development stages: pre-anthesis, anthesis and post-anthesis.

139 For the characterization experiment of flower events, a description of flower habits was
140 performed and compared with those available in the literature. All the results were recorded
141 through photography and expressed as percentage.

142 For the experiments of pollen viability and stigma receptivity, the data were submitted to an
143 analysis of variance, being the pollen viability experiment subjected to linear or quadratic
144 regression at 5% probability level. The stigma receptivity experiment was subjected to Scott-
145 Knott test at 5% error probability. All of the analyses were performed on the Sisvar® statistical
146 software (FERREIRA, 2011).

147 **RESULTS AND DISCUSSION**

148 During the evaluation of flower events, it was verified that all the flower buds of the
149 inflorescences did not show a synchronous opening (lower than 6%), thus observing flowers at
150 different stages in the same inflorescence (Figures 1A, B and C).

151 **Figure 1** - Inflorescences of olive tree (*Olea europaea* L.), Arbequina cultivar. (A) Onset of
 152 flowering, showing pre-anthesis flowers with different sizes and flowers at the beginning of the
 153 anthesis - detail of non-dehiscent anthers (arrows). (B) Inflorescence containing flowers in pre-
 154 anthesis, anthesis and post-anthesis in the same inflorescence. (C) Post-anthesis of all flowers of
 155 an inflorescence - detail for the senescence of the corolla (black arrow), flowers without corolla,
 156 with stigma showing an intense yellow coloration (red arrow). Bar: 5 mm



157

158

159 This information goes against the strategy of several fruit species in temperate regions,
 160 where the plants come into dormancy in autumn and winter, showing a fast and simultaneous
 161 flowering during the spring, guaranteeing quick pollination and fertilization, which causes
 162 homogeneity in the fruit development (GUO *et al.*, 2014).

163 However, such non-uniformity in olive flowering is intrinsic to the species, similarly as
 164 other phenological records found in the literature (OLIVEIRA, M., *et al.*, 2012; SADOK *et al.*,
 165 2013; SANZ-CORTÉS *et al.*, 2002). This fact is probably due to the olive tree diminishing the
 166 metabolic activity, but without coming into complete dormancy in the colder seasons, which
 167 affects the synchrony during the differentiation of flower buds in the spring. Furthermore,
 168 thermal oscillations during the winter period may lead to the physiological decompensation of
 169 these buds, specifically related to the chill accumulation, also causing uniformity and reduction in
 170 flowering (GUO *et al.*, 2014).

171 Also during the evaluation, it was observed that in the pre-anthesis (closed bud), there was
172 no release of pollen grains on the stigma; all the flowers showed indehiscent anthers at the time
173 of flower opening (Figure 1A), refuting the possibility of occurrence of cleistogamy in flowers.


174 Flower opening occurs more frequently between 10 a.m. and 11 p.m. However, the anthers
175 are not dehiscent in 82% of the evaluated flowers; its desiccation and release of pollen grain
176 occur gradually throughout two or more days in 80% of the flowers (Figure 1A). This
177 characteristic of gradual release of pollen grains throughout the days is sharper in plants with
178 anemophily, in which the temperature and relative humidity directly influence the anther
179 desiccation and release of the pollen grain (MARTIN; CHAMECKI; BRUSH, 2010).

180 Flower longevity (period between anthesis and corolla senescence) in cv. Arbequina was
181 approximately one week in 84% of the analyzed flowers. This information also corroborates with
182 Suárez, Castro and Rapoport (2012), which worked with morphological and histochemical
183 alterations of the pistil of olive trees and found during their phenological evaluation that it can
184 take more than a week between the onset of anthesis and its senescence.

185 In the pollen grain conservation experiment, it is verified through linear regression (Figure
186 2) that the pollen viability was drastically decreased at room temperature after the second day of
187 storage (close to 5%), being that the material no longer responded to the germination test in
188 culture medium after 40 days.

189
190
191
192
193

194 **Figure 2** - Pollen viability of olive grains (*Olea europaea* L.), Arbequina cultivar, stored for 105
 195 days at different storage locations (room temperature at 27 °C, refrigerator at 8 °C and freezer at -
 196 10 °C



197
 198
 199 For several studies with fruit trees, 30% germination of pollen grains is considered an
 200 acceptable rate to ensure good fertilization and fruiting (ALBUQUERQUE JÚNIOR, 2010;
 201 SILVA *et al.*, 2016). Considering the germination rate around 5% (Figure 2), it can be considered
 202 as very low for the use of this material in hybridization studies in olive trees.

203 Regarding the other treatments, the storage of pollen grains at 8 °C showed a viability
 204 around 56%, being that there was a drastic decrease in the pollen germination after that period,
 205 becoming unviable after 63 days (Figure 2).

206 The freezer storage method at -10 °C showed higher results in relation to the other
 207 treatments, preserving the material viability for longer time (Figure 2). At 60 days of storage, the
 208 pollen grains still had a germination rate around 30%, i.e., an acceptable parameter for the use,
 209 according to the literature (ALBUQUERQUE JÚNIOR, 2010; HAUAGGE; BRUCKNER,
 210 2002). Fifteen days after this period, the material still showed a low germination rate when

211 subjected to the culture medium, becoming completely unviable only 99 days after the
212 experiment.


213 With respect to the material viability, it is possible to observe that the pollen grains stored
214 at 27 °C (ambient temperature) lost viability quickly. This fact may be related to the high
215 temperature of the environment, which maintains or accelerates the material's metabolic activity,
216 making it unviable quickly. The storage at ambient temperature also makes the material more
217 susceptible to moisture absorption, which accelerates the metabolism, besides making pollen
218 grains more susceptible to attack by fungi and microorganisms (KOUBOURIS; METZIDAKIS;
219 VASILAKAKIS, 2009).

220 It was also possible to observe that treatments with lower temperatures showed greater
221 durability. The use of low temperatures is related to the reduction of the metabolism of pollen
222 grains, which favors greater longevity of the material (CUCHIARA; SILVA; BOBROWSHI,
223 2012).

224 Regarding the stigma receptivity evaluation, it was verified that there was no statistical
225 difference among the treatments; the stigmas of all development stages and at all times show
226 bubble formation in the stigma cavity when subjected to the 3% hydrogen peroxide test,
227 indicating activity of the peroxidase enzyme (Figure 2).

228
229
230
231
232
233
234
235

236 **Figure 3** - Flower development of olive tree (*Olea europaea* L.), Arbequina cultivar, and stigma
 237 receptivity at different development stages, using the 3% hydrogen peroxide (H₂O₂) method. (A
 238 and B) Pre-anthesis flowers. (C and D) Anthesis flowers. (E and F) Post-anthesis flowers. (G)
 239 Extremely green flower bud, pre-anthesis, showing receptive stigma. (H) Flower bud stigma,
 240 fully developed, pre-anthesis and receptive. (I) Anthesis flower, detail for receptive stigma cavity
 241 (arrow). (J) Post-anthesis flower with receptive stigma. Bar: 1mm



242
 243
 244 These results are highly relevant for breeding experiments. Once the pre-anthesis flowers
 245 are receptive, controlled hybridization can be performed without contamination of pollen grains
 246 other than the selected material (Figures 3A, 3B, 3G and 3H).

247 However, it is recommended to the breeder the choice of developed flower buds and as
 248 close as possible to the anthesis (Figures 3B and 3H), since although the stigma of olive tree is
 249 receptive, it is only close to the flower anthesis that the style transmission will begin to break the
 250 starch and hence provide nutrients for the development of the pollen tube (SUÀREZ; CASTRO;
 251 RAPOPORT, 2012).

252 Regarding the time of an open flower, the duration of gynoecium receptivity will be
 253 directly related to the supply of pollen grain in the environment (LANKINEN; MADJIDIAN,
 254 2011). In other words, the higher the pollen grain supply, the lower the time of stigma reception,
 255 the lower the pollen supply, the longer the time of its receptivity.

256 Additionally, several studies with different species have shown that flower longevity
 257 increases according to the altitude. This basically occurs due to the lower pollination rates, and/or
 258 due to the lower temperatures that lead to decreased metabolic activity, delaying the anthesis
 259 (MU *et al.*, 2011; PACHECO *et al.*, 2016; PÉLABON *et al.*, 2013; STEINACHER; WAGNER,
 260 2010), which is also a possible explanation for the receptivity period of this experiment.

261 **CONCLUSIONS**

- 262 1. The olive pollen grains, Arbequina cultivar, are preserved as viable for use in hybridizations
 263 for up to 60 days when stored in the freezer at -10 °C;
- 264 2. The stigmas are receptive since pre-anthesis, with receptivity frequency prevailing throughout
 265 the day, allowing performing hybridizations at different flower development stages and at
 266 different daytimes.

267 **ACKNOWLEDGEMENTS**

268 The authors would like to thank the Foundation for Supporting Research of the State of
 269 Minas Gerais (FAPEMIG), the National Council for Scientific and Technological Development
 270 (CNPq) and the Coordination for the Improvement of Higher Education Personnel (CAPES) for
 271 the financial support to perform this study.

272 **REFERENCES**

- 273 ALBUQUERQUE JÚNIOR, C. L. *et al.* Número de anteras por flor, grãos de pólen por antera e
 274 capacidade germinativa do pólen de diferentes cultivares de macieiras. **Revista Brasileira de**
 275 **Fruticultura**, v. 32, n. 4, p. 1255-1260, 2010.

- 276 BRITO, A. C. *et al.* Propriedade do pólen e do stigma de *Ocimum basilicum* L. (cultivar Maria
277 Bonita) para aumentar a eficiência de cruzamentos em programas de melhoramento. **Revista**
278 **Brasileira de Plantas Medicinais**, v. 12, n. 2, p. 208-214, 2010.
- 279 CONSEJO OLEÍCOLA INTERNACIONAL. **Cifras aceites de oliva en aceitonas de mesa.**
280 2017. Disponível em: <[http://www.internationaloliveoil.org/estaticos/view/131-world-olive-oil-](http://www.internationaloliveoil.org/estaticos/view/131-world-olive-oil-figures)
281 [figures](http://www.internationaloliveoil.org/estaticos/view/131-world-olive-oil-figures)>. Acesso em: 31 maio 2017.
- 282 CUCHIARA, C. C.; SILVA, S. D. A.; BOBROWSKI, V. L. Conservação de grãos de pólen de
283 mamoneiro a baixas temperaturas. **Revista Ceres**, v. 59, n. 1, p. 82-87, 2012.
- 284 FERREIRA, D. F. Sisvar: a computer statistical analysis system. **Ciência e Agrotecnologia**, v.
285 35, n. 6, p. 1039-1042, 2011.
- 286 FIGUEIREDO, M. A. *et al.* Características florais e carpométricas e germinação in vitro de grãos
287 de pólen de cultivares de amoreira-preta. **Pesquisa Agropecuária Brasileira**, v. 48, n. 7, p. 731-
288 740, 2013.
- 289 GUO, L. *et al.* Chilling and requirements for flowering in temperate fruit trees. **International**
290 **Journal of Biometeorology**, v. 58, n. 6, p. 1195-1206, 2014.
- 291 HAUAGGE, R.; BRUCKNER, C. H. **Melhoramento de fruteiras de clima temperado.** Viçosa,
292 MG: UFV, 2002. 88 p.
- 293 KEARNS, C. A.; INOUE, D. W. **Techniques for pollination biologist.** Niwot: University
294 Press of Colorado, 1993. 579 p.
- 295 KOUBOURIS, G. C.; METZIDAKIS, I. T.; VASILAKAKIS, M. D. Impact of temperature on
296 olive (*Olea europaea* L.) pollen performance in relation to relative humidity and genotype.
297 **Environmental and Experimental Botany**, v. 67, n. 1, p. 209-214, 2009.

- 298 LANKINEN, A.; MADJIDIAN, J. A.; Enhancing pollen competition by delayaing stigma
299 receptivity: pollen deposition schedules affect sering ability, paternal diversity, and seed
300 production in *Collinsia heterophylla* (Pantaginaceae). **American Journal of Botany**, v. 98, n. 7,
301 p. 1191-1200, 2011.
- 302 MARTIN, M. D.; CHAMECKI, M.; BRUSH, G. Anthesis synchronization and floral morphology
303 determine diurnal patterns of ragweed pollen dispersal. **Agricultural and Forest Meteorology**,
304 v. 150, n. 9, p. 1307-1317, 2010.
- 305 MOURA, C. R. F.; MACHADO, C. A.; LÈDO, A. S. In vitro germination and viability of pollen
306 grain of coconut accessions. **Revista Ciência Agronomica**, v. 46, n. 2, p. 421-427, 2015.
- 307 MU, J. *et al.* Difference in floral traits, pollination, and reproductive success between white and
308 blue flowers of *Gentiana leucomelaena* (Gentianaceae) in na alpine meadow. **Artic, Antartic,**
309 **and Alpine Research**, v. 43, n. 3, p. 410-416, 2011.
- 310 OLIVEIRA, A. F. *et al.* Caracterização e proteção de cultivares. *In:* OLIVEIRA, A. F. **Oliveiras**
311 **do Brasil tecnologia de produção**. 1. ed. Belo Horizonte: Epamig, 2012. cap. 9, p. 251-274.
- 312 OLIVEIRA, M. C. *et al.* Características fenológicas e físicas e perfil de ácidos graxos em
313 oliveiras no sul de Minas Gerais. **Pesquisa Agropecuária Brasileira**, v. 47, n. 1, p. 30-35, 2012.
- 314 PACHECO, D. A. *et al.* Plastic respinses contribute to explaining altitudinal and temporal
315 variation in potential flower longevity in high andrean *Rhodolirion montanum*. **Plos One**, v. 11,
316 n. 11, p. 1-18, 2016.
- 317 PÉLABON, C. *et al.* Decoupled phenotypic variation between floral and vegetative traits:
318 distinguishing between developmental and environmental correlations. **Annals of Botany**, v.
319 111, n. 5, p. 935-944, 2013.

- 320 SADOK, I. B. *et al.* QTL mapping of flowering and fruting traits in olive. **Plos One**, v. 8, n. 5, p.
321 1-16, 2013.
- 322 SANZ-CORTÉS, F. *et al.* Phenological growth stages of olive tress (*Olea europaea*). **Annals of**
323 **Applied Biology**, v. 140, n. 2, p. 151-157, 2002.
- 324 SCHIFINO-WITTMANN, M. T.; DALL'AGNOL, M. Auto-Incompatibilidade em plantas.
325 **Ciência Rural**, v. 32, n. 6, p. 1083-1090, 2002.
- 326 SILVA, L. F. O. *et al.* Caracterização agronômica e carpométrica de cultivares de oliveira.
327 **Pesquisa Agropecuária Tropical**, v. 42, n. 3, p. 350-356, 2012b.
- 328 SILVA, L. F. O. *et al.* Establishment of growth medium and quantification of pollen grains of
329 olive cultivars in Brazil's subtropical areas. **Bragantia**, v. 75, n. 1, p. 26-32, 2016.
- 330 SILVA, L. F. O. *et al.* Variação na qualidade do azeite em cultivares de oliveira. **Bragantia**, v.
331 71, n. 2, p. 202-209, 2012a.
- 332 STEINACHER, G.; WAGNER, J. Flower longevity and of pistil receptivity in high mountain
333 plants. **Flora**, v. 205, n. 6, p. 376-387, 2010.
- 334 SUÀREZ, C.; CASTRO, A. J.; RAPOPORT, H. F. Morphological, histological and
335 ultrastructural changes in the olive pistil during flowering. **Sexual Plant Reproduction**, v. 25, n.
336 2, p. 133-146, 2012.